

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Agriculture and Natural Resources

6-25-2007

SCN females on roots signal infestations and possibly ineffectiveness of resistance

Gregory L. Tylka

Iowa State University, gltylka@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Plant Pathology Commons](#)

Recommended Citation

Tylka, Gregory L., "SCN females on roots signal infestations and possibly ineffectiveness of resistance" (2007). *Integrated Crop Management News*. Paper 1038.

<http://lib.dr.iastate.edu/cropnews/1038>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact digirep@iastate.edu.

INTEGRATED CROP MANAGEMENT

Get the latest research-based information on crops. [Sign up to be notified](#) when new content is available!

ICM > 2007 > IC-498(16) -- June 25, 2007

Current Newsletter

You are viewing **archives** for the newsletter from 1993-2007. For current news, see [Integrated Crop Management News](#).

Archives 1993-2007



Announcements



Crop Production



Insects and Mites



Pesticide Education



Plant Diseases



Soils



Weed Management

Image Gallery

Printable Version

Printable version of this page

Related Articles

Surveying Iowa for SCN
December 10, 2007

Field testing of N-Hibit[®]; seed treatment in 2007 for management of SCN in Iowa

December 10, 2007

A new take on soil sampling fields for SCN
November 12, 2007

SCN females on roots signal infestations and possibly ineffectiveness of resistance

by Greg Tylka, Department of Plant Pathology

Many fields are not known to be infested with the soybean cyst nematode (SCN) because obvious, above-ground symptoms do not always occur. A reliable way to check for SCN on non-resistant (susceptible) soybean varieties is to dig roots, carefully crumble away much of the soil from the roots, and look for the adult SCN females. The females appear as small, round, white objects on the roots and are about the size of a period at the end of a sentence.

In addition to checking fields for SCN infestations, looking for SCN females on roots of SCN-resistant soybeans can serve as a check of the effectiveness of resistant varieties in controlling SCN reproduction.

Resistance to SCN in soybeans is not 100 percent effective, so it is normal to see a few (up to 15 or so) SCN females on the roots of resistant soybeans. But resistance may be failing if many (100 or more) SCN females are seen on resistant soybean roots. In 2006, there were numerous reports of many SCN females being seen on roots of SCN-resistant soybean varieties in Iowa.

A survey is currently underway, funded by the Iowa Soybean Association, to determine how well SCN populations from throughout Iowa can reproduce on SCN-resistant soybean varieties.

ISU Extension field agronomist John Holmes observed adult SCN females on soybean roots near Webster City in Hamilton County in mid-June 2007. The soybeans were planted in early May, and the SCN females were on roots of plants growing in high pH spots. SCN females should be apparent on young roots of soybean plants in Iowa through July and August. Roots should be checked no earlier than 4 or 5 weeks after planting.

Additional information about the biology, sampling, and management of SCN can be found on the Web at www.soybeancyst.info.

Results of SCN-resistant soybean variety testing become available

November 12, 2007

Sample fields this fall for SCN to figure out 2007 or plan for 2008

October 1, 2007

Winter annual weeds and SCN: What's the connection?

April 23, 2007

Soybean cyst nematode: Still a major threat to soybean production

March 26, 2007

SCN-resistant soybean varieties: Not all are created equal

February 26, 2007

How to interpret SCN soil test results

December 18, 2006

What's your type?: An HG type test for SCN populations

November 13, 2006



SCN-resistant soybeans damaged by SCN in southwest Iowa. (Greg Tylka)



Adult soybean cyst nematode females on soybean roots at tip of key. (Greg Tylka)

Greg Tylka is a professor of plant pathology with extension and research responsibilities in management of plant-parasitic nematodes.

This article originally appeared on page 192 of the IC-498(16) -- June 25, 2007 issue.

Updated 06/28/2007 - 1:57pm

